

What generic automorphisms of the random poset look like

Dakota Thor Ihli
University of Illinois at Urbana-Champaign

Online Logic Seminar
4 Mar 2021

What are “generic automorphisms”?

What are “generic automorphisms”?

Definition

For a Polish group G , an element $g \in G$ is called **generic** if it lies in a (necessarily unique) comeagre conjugacy class.

What are “generic automorphisms”?

Definition

For a Polish group G , an element $g \in G$ is called **generic** if it lies in a (necessarily unique) comeagre conjugacy class.

If \mathbf{K} is a countable (first-order) structure, its automorphism group $\text{Aut}(\mathbf{K})$ is a Polish group with the pointwise convergence topology.

What are “generic automorphisms”?

Definition

For a Polish group G , an element $g \in G$ is called **generic** if it lies in a (necessarily unique) comeagre conjugacy class.

If \mathbf{K} is a countable (first-order) structure, its automorphism group $\text{Aut}(\mathbf{K})$ is a Polish group with the pointwise convergence topology.

Thus, a **generic automorphism** of \mathbf{K} is a generic element of $\text{Aut}(\mathbf{K})$.

What structures do we care about here?

What structures do we care about here?

Definition

Fix a relational language L , and let \mathbf{K} be a countable structure.

What structures do we care about here?

Definition

Fix a relational language L , and let \mathbf{K} be a countable structure.

- A **partial automorphism** of \mathbf{K} is an isomorphism between substructures of \mathbf{K} .

What structures do we care about here?

Definition

Fix a relational language L , and let \mathbf{K} be a countable structure.

- A **partial automorphism** of \mathbf{K} is an isomorphism between substructures of \mathbf{K} . Let $\text{Aut}_{<\omega}(\mathbf{K})$ denote the set of finite partial automorphisms of \mathbf{K} .

What structures do we care about here?

Definition

Fix a relational language L , and let \mathbf{K} be a countable structure.

- A **partial automorphism** of \mathbf{K} is an isomorphism between substructures of \mathbf{K} . Let $\text{Aut}_{<\omega}(\mathbf{K})$ denote the set of finite partial automorphisms of \mathbf{K} . For each $p \in \text{Aut}_{<\omega}(\mathbf{K})$, let $[p]$ denote the set of $f \in \text{Aut}(\mathbf{K})$ extending p .

What structures do we care about here?

Definition

Fix a relational language L , and let \mathbf{K} be a countable structure.

- A **partial automorphism** of \mathbf{K} is an isomorphism between substructures of \mathbf{K} . Let $\text{Aut}_{<\omega}(\mathbf{K})$ denote the set of finite partial automorphisms of \mathbf{K} . For each $p \in \text{Aut}_{<\omega}(\mathbf{K})$, let $[p]$ denote the set of $f \in \text{Aut}(\mathbf{K})$ extending p .
- \mathbf{K} is **ultrahomogeneous** if every finite partial automorphism of \mathbf{K} extends to a (full) automorphism of \mathbf{K} .

What structures do we care about here?

Definition

Fix a relational language L , and let \mathbf{K} be a countable structure.

- A **partial automorphism** of \mathbf{K} is an isomorphism between substructures of \mathbf{K} . Let $\text{Aut}_{<\omega}(\mathbf{K})$ denote the set of finite partial automorphisms of \mathbf{K} . For each $p \in \text{Aut}_{<\omega}(\mathbf{K})$, let $[p]$ denote the set of $f \in \text{Aut}(\mathbf{K})$ extending p .
- \mathbf{K} is **ultrahomogeneous** if every finite partial automorphism of \mathbf{K} extends to a (full) automorphism of \mathbf{K} . Equivalently, if $[p]$ is non-empty for all $p \in \text{Aut}_{<\omega}(\mathbf{K})$.

What structures do we care about here?

Definition

Fix a relational language L , and let \mathbf{K} be a countable structure.

- A **partial automorphism** of \mathbf{K} is an isomorphism between substructures of \mathbf{K} . Let $\text{Aut}_{<\omega}(\mathbf{K})$ denote the set of finite partial automorphisms of \mathbf{K} . For each $p \in \text{Aut}_{<\omega}(\mathbf{K})$, let $[p]$ denote the set of $f \in \text{Aut}(\mathbf{K})$ extending p .
- \mathbf{K} is **ultrahomogeneous** if every finite partial automorphism of \mathbf{K} extends to a (full) automorphism of \mathbf{K} . Equivalently, if $[p]$ is non-empty for all $p \in \text{Aut}_{<\omega}(\mathbf{K})$.
- If \mathfrak{K} is a class of L -structures, \mathbf{K} is **universal** for \mathfrak{K} if every structure in \mathfrak{K} embeds into \mathbf{K} .

What structures do we care about here?

We are interested in **Fraïssé structures**: ultrahomogeneous countable structures which are universal for a coherent class of finite structures called a **Fraïssé class**.

What structures do we care about here?

We are interested in **Fraïssé structures**: ultrahomogeneous countable structures which are universal for a coherent class of finite structures called a **Fraïssé class**.

Examples

Fraïssé class	Fraïssé structure
---------------	-------------------

What structures do we care about here?

We are interested in **Fraïssé structures**: ultrahomogeneous countable structures which are universal for a coherent class of finite structures called a **Fraïssé class**.

Examples

Fraïssé class	Fraïssé structure
{finite sets}	

What structures do we care about here?

We are interested in **Fraïssé structures**: ultrahomogeneous countable structures which are universal for a coherent class of finite structures called a **Fraïssé class**.

Examples

Fraïssé class	Fraïssé structure
{finite sets}	ω

What structures do we care about here?

We are interested in **Fraïssé structures**: ultrahomogeneous countable structures which are universal for a coherent class of finite structures called a **Fraïssé class**.

Examples

Fraïssé class	Fraïssé structure
{finite sets}	ω
{finite linear orders}	

What structures do we care about here?

We are interested in **Fraïssé structures**: ultrahomogeneous countable structures which are universal for a coherent class of finite structures called a **Fraïssé class**.

Examples

Fraïssé class	Fraïssé structure
{finite sets}	ω
{finite linear orders}	$(\mathbb{Q}, <)$

What structures do we care about here?

We are interested in **Fraïssé structures**: ultrahomogeneous countable structures which are universal for a coherent class of finite structures called a **Fraïssé class**.

Examples

Fraïssé class	Fraïssé structure
{finite sets}	ω
{finite linear orders}	$(\mathbb{Q}, <)$
{finite (undirected) graphs}	

What structures do we care about here?

We are interested in **Fraïssé structures**: ultrahomogeneous countable structures which are universal for a coherent class of finite structures called a **Fraïssé class**.

Examples

Fraïssé class	Fraïssé structure
{finite sets}	ω
{finite linear orders}	$(\mathbb{Q}, <)$
{finite (undirected) graphs}	The random graph

What structures do we care about here?

We are interested in **Fraïssé structures**: ultrahomogeneous countable structures which are universal for a coherent class of finite structures called a **Fraïssé class**.

Examples

Fraïssé class	Fraïssé structure
{finite sets}	ω
{finite linear orders}	$(\mathbb{Q}, <)$
{finite (undirected) graphs}	The random graph
{finite partial orders}	

What structures do we care about here?

We are interested in **Fraïssé structures**: ultrahomogeneous countable structures which are universal for a coherent class of finite structures called a **Fraïssé class**.

Examples

Fraïssé class	Fraïssé structure
{finite sets}	ω
{finite linear orders}	$(\mathbb{Q}, <)$
{finite (undirected) graphs}	The random graph
{finite partial orders}	The random poset

What structures do we care about here?

We are interested in **Fraïssé structures**: ultrahomogeneous countable structures which are universal for a coherent class of finite structures called a **Fraïssé class**.

Examples

Fraïssé class	Fraïssé structure
{finite sets}	ω
{finite linear orders}	$(\mathbb{Q}, <)$
{finite (undirected) graphs}	The random graph
{finite partial orders}	The random poset =: P

Their automorphism groups:

Their automorphism groups:

Remark

All the examples on the previous slide admit generic automorphisms. (That is, their automorphism groups have comeagre conjugacy classes.)

Their automorphism groups:

Remark

All the examples on the previous slide admit generic automorphisms. (That is, their automorphism groups have comeagre conjugacy classes.)

What might generic automorphisms *look like*? For example:

Theorem (Truss, 1991)

Let $f \in S_\infty$.

Their automorphism groups:

Remark

All the examples on the previous slide admit generic automorphisms. (That is, their automorphism groups have comeagre conjugacy classes.)

What might generic automorphisms *look like*? For example:

Theorem (Truss, 1991)

Let $f \in S_\infty$. Then f is generic if and only if:

Their automorphism groups:

Remark

All the examples on the previous slide admit generic automorphisms. (That is, their automorphism groups have comeagre conjugacy classes.)

What might generic automorphisms *look like*? For example:

Theorem (Truss, 1991)

Let $f \in S_\infty$. Then f is generic if and only if:

- f has no infinite orbits;

Their automorphism groups:

Remark

All the examples on the previous slide admit generic automorphisms. (That is, their automorphism groups have comeagre conjugacy classes.)

What might generic automorphisms *look like*? For example:

Theorem (Truss, 1991)

Let $f \in S_\infty$. Then f is generic if and only if:

- f has no infinite orbits;
- For every n , f has infinitely many orbits of length n .

Their automorphism groups:

Remark

All the examples on the previous slide admit generic automorphisms. (That is, their automorphism groups have comeagre conjugacy classes.)

What might generic automorphisms *look like*? For example:

Theorem (Truss, 1991)

Let $f \in S_\infty$. Then f is generic if and only if:

- *f has no infinite orbits;*
- *For every n , f has infinitely many orbits of length n .*

Remark

Generic automorphisms of $(\mathbb{Q}, <)$ and the random graph admit similar kinds of descriptions.

What about the random poset?

Theorem (Kuske–Truss, 2000)

The random poset \mathbf{P} admits generic automorphisms.

What about the random poset?

Theorem (Kuske–Truss, 2000)

The random poset \mathbf{P} admits generic automorphisms.

Goal

Find an explicit description of generic automorphisms of \mathbf{P} .

Tools for describing automorphisms' actions

Tools for describing automorphisms' actions

Definition

Let $(P, <)$ be any poset, and let $f \in \text{Aut}(P)$.

Tools for describing automorphisms' actions

Definition

Let $(P, <)$ be any poset, and let $f \in \text{Aut}(P)$.

- The **spiral length** of x , denoted $\text{sp}(x, f)$,

Tools for describing automorphisms' actions

Definition

Let $(P, <)$ be any poset, and let $f \in \text{Aut}(P)$.

- The **spiral length** of x , denoted $\text{sp}(x, f)$, is the least $n \geq 1$ for which x and $f^n(x)$ are comparable,

Tools for describing automorphisms' actions

Definition

Let $(P, <)$ be any poset, and let $f \in \text{Aut}(P)$.

- The **spiral length** of x , denoted $\text{sp}(x, f)$, is the least $n \geq 1$ for which x and $f^n(x)$ are comparable, or ∞ if no such n exists.

Tools for describing automorphisms' actions

Definition

Let $(P, <)$ be any poset, and let $f \in \text{Aut}(P)$.

- The **spiral length** of x , denoted $\text{sp}(x, f)$, is the least $n \geq 1$ for which x and $f^n(x)$ are comparable, or ∞ if no such n exists.
- The **parity** of x

Tools for describing automorphisms' actions

Definition

Let $(P, <)$ be any poset, and let $f \in \text{Aut}(P)$.

- The **spiral length** of x , denoted $\text{sp}(x, f)$, is the least $n \geq 1$ for which x and $f^n(x)$ are comparable, or ∞ if no such n exists.
- The **parity** of x is given by:


$$\text{par}(x, f) := \begin{cases} +1 & \text{if } \text{sp}(x, f) = n < \infty \text{ and } x < f^n(x); \\ -1 & \text{if } \text{sp}(x, f) = n < \infty \text{ and } x > f^n(x); \\ 0 & \text{otherwise.} \end{cases}$$

Tools for describing automorphisms' actions

$$\begin{array}{c} f^3(x) \\ | \\ x \end{array} \quad f(x) \quad f^2(x)$$

Example: suppose $\text{sp}(x, f) = 3$
and $\text{par}(x, f) = +1$.


Tools for describing automorphisms' actions



Example: suppose $\text{sp}(x, f) = 3$
and $\text{par}(x, f) = +1$.

Since f is an automorphism,
 $f^k(x) < f^{k+3}(x)$ for all $k \in \mathbb{Z}$.


Tools for describing automorphisms' actions



Example: suppose $\text{sp}(x, f) = 3$
and $\text{par}(x, f) = +1$.

Since f is an automorphism,
 $f^k(x) < f^{k+3}(x)$ for all $k \in \mathbb{Z}$.
This breaks the orbit $f^{\mathbb{Z}}(x)$ into
“rails”,


Tools for describing automorphisms' actions



Example: suppose $\text{sp}(x, f) = 3$
and $\text{par}(x, f) = +1$.

Since f is an automorphism,
 $f^k(x) < f^{k+3}(x)$ for all $k \in \mathbb{Z}$.
This breaks the orbit $f^{\mathbb{Z}}(x)$ into
"rails", but there may be other
relations too.

Tools for describing automorphisms' actions



Example: suppose $\text{sp}(x, f) = 3$
and $\text{par}(x, f) = +1$.

Since f is an automorphism,
 $f^k(x) < f^{k+3}(x)$ for all $k \in \mathbb{Z}$.
This breaks the orbit $f^{\mathbb{Z}}(x)$ into
“rails”, but there may be other
relations too.

More tools for describing automorphisms' actions

Definition

Let $(P, <)$ be any poset, and let $f \in \text{Aut}(P)$.

More tools for describing automorphisms' actions

Definition

Let $(P, <)$ be any poset, and let $f \in \text{Aut}(P)$.

- Define $x \sim_f y \stackrel{\text{def}}{\iff} \exists i, j \in \mathbb{Z} (f^i(x) \leq y \leq f^j(x))$.

More tools for describing automorphisms' actions

Definition

Let $(P, <)$ be any poset, and let $f \in \text{Aut}(P)$.

- Define $x \sim_f y \stackrel{\text{def}}{\iff} \exists i, j \in \mathbb{Z} (f^i(x) \leq y \leq f^j(x))$.
- \sim_f is an equivalence relation.

More tools for describing automorphisms' actions

Definition

Let $(P, <)$ be any poset, and let $f \in \text{Aut}(P)$.

- Define $x \sim_f y \stackrel{\text{def}}{\iff} \exists i, j \in \mathbb{Z} (f^i(x) \leq y \leq f^j(x))$.
- \sim_f is an equivalence relation. Let the quotient map be denoted by $\mathcal{O}_f : P \rightarrow P / \sim_f$.

More tools for describing automorphisms' actions

Definition

Let $(P, <)$ be any poset, and let $f \in \text{Aut}(P)$.

- Define $x \sim_f y \stackrel{\text{def}}{\iff} \exists i, j \in \mathbb{Z} (f^i(x) \leq y \leq f^j(x))$.
- \sim_f is an equivalence relation. Let the quotient map be denoted by $\mathcal{O}_f : P \rightarrow P / \sim_f$.
- The equivalence classes $\mathcal{O}_f(x)$ are called **orbitals**.


More tools for describing automorphisms' actions

Definition


Let $(P, <)$ be any poset, and let $f \in \text{Aut}(P)$.

- Define $x \sim_f y \stackrel{\text{def}}{\iff} \exists i, j \in \mathbb{Z} (f^i(x) \leq y \leq f^j(x))$.
- \sim_f is an equivalence relation. Let the quotient map be denoted by $\mathcal{O}_f : P \rightarrow P / \sim_f$.
- The equivalence classes $\mathcal{O}_f(x)$ are called **orbitals**.
- The quotient $\mathcal{O}_f[P]$ is called the **orbital quotient**.

More tools for describing automorphisms' actions



More tools for describing automorphisms' actions



Orbitals are the “convex hulls” of orbits.

More tools for describing automorphisms' actions

Facts

More tools for describing automorphisms' actions

Facts

- $f^{\mathbb{Z}}(x) \subseteq \mathcal{O}_f(x)$, and equality holds whenever $\text{par}(x, f) = 0$.

More tools for describing automorphisms' actions

Facts

- $f^{\mathbb{Z}}(x) \subseteq \mathcal{O}_f(x)$, and equality holds whenever $\text{par}(x, f) = 0$.
- Parity is orbital-invariant; that is, $x \sim_f y$ implies $\text{par}(x, f) = \text{par}(y, f)$.

More tools for describing automorphisms' actions

Facts

- $f^{\mathbb{Z}}(x) \subseteq \mathcal{O}_f(x)$, and equality holds whenever $\text{par}(x, f) = 0$.
- Parity is orbital-invariant; that is, $x \sim_f y$ implies $\text{par}(x, f) = \text{par}(y, f)$.
- Spiral length *need not be* orbital-invariant (unless $\text{par}(x, f) = 0$).

Example in \mathbb{Q} :

$$\text{Let } f(x) := \begin{cases} 2x + 3 & x \leq -2, \\ \frac{x}{2} & -2 \leq x \leq 2, \\ 2x - 3 & 2 \leq x. \end{cases}$$

Example in \mathbb{Q} :

$$\text{Let } f(x) := \begin{cases} 2x + 3 & x \leq -2, \\ \frac{x}{2} & -2 \leq x \leq 2, \\ 2x - 3 & 2 \leq x. \end{cases}$$



Then there are seven orbitals: three of parity 0, two each of parity -1 and $+1$.

Orbitals inherit order structure from \mathbf{P} !

Definition

Define two orders on $\mathcal{O}_f[P]$ — one strong and one weak:

Orbitals inherit order structure from **P**!

Definition

Define two orders on $\mathcal{O}_f[P]$ — one **strong** and one weak:

$$\mathcal{O}_f(x) <_f^s \mathcal{O}_f(y) \stackrel{\text{def}}{\iff} x' < y' \text{ for all } x' \sim_f x \text{ and } y' \sim_f y,$$

Orbitals inherit order structure from \mathbf{P} !

Definition

Define two orders on $\mathcal{O}_f[P]$ — one **strong** and one **weak**:

$$\mathcal{O}_f(x) <_f^s \mathcal{O}_f(y) \stackrel{\text{def}}{\iff} x' < y' \text{ for } \mathbf{all} \ x' \sim_f x \text{ and } y' \sim_f y,$$

$$\mathcal{O}_f(x) \leq_f^w \mathcal{O}_f(y) \stackrel{\text{def}}{\iff} x' \leq y' \text{ for } \mathbf{some} \ x' \sim_f x \text{ and } y' \sim_f y.$$

Orbitals inherit order structure from \mathbf{P} !

Definition

Define two orders on $\mathcal{O}_f[P]$ — one **strong** and one **weak**:

$$\mathcal{O}_f(x) <_f^s \mathcal{O}_f(y) \stackrel{\text{def}}{\iff} x' < y' \text{ for } \mathbf{all} \ x' \sim_f x \text{ and } y' \sim_f y,$$

$$\mathcal{O}_f(x) \leq_f^w \mathcal{O}_f(y) \stackrel{\text{def}}{\iff} x' \leq y' \text{ for } \mathbf{some} \ x' \sim_f x \text{ and } y' \sim_f y.$$

Remark

If P is linearly ordered, these orders agree and are linear orders themselves.

Using orbitals to characterize generics

Using orbitals to characterize generics

Theorem (Truss, 1991)

Let $f \in \text{Aut}(\mathbb{Q})$.

Using orbitals to characterize generics

Theorem (Truss, 1991)

Let $f \in \text{Aut}(\mathbb{Q})$. Then f is generic if and only if $\mathcal{O}_f[\mathbb{Q}] \cong \mathbb{Q}$,

Using orbitals to characterize generics

Theorem (Truss, 1991)

Let $f \in \text{Aut}(\mathbb{Q})$. Then f is generic if and only if $\mathcal{O}_f[\mathbb{Q}] \cong \mathbb{Q}$, and for each $\sigma \in \{+1, -1, 0\}$, the set of orbitals of parity σ is dense in $\mathcal{O}_f[\mathbb{Q}]$.

Using orbitals to characterize generics

Theorem (Truss, 1991)

Let $f \in \text{Aut}(\mathbb{Q})$. Then f is generic if and only if $\mathcal{O}_f[\mathbb{Q}] \cong \mathbb{Q}$, and for each $\sigma \in \{+1, -1, 0\}$, the set of orbitals of parity σ is dense in $\mathcal{O}_f[\mathbb{Q}]$.

Question

Is an analogous statement true for **P**?

Using orbitals to characterize generics

Theorem (Truss, 1991)

Let $f \in \text{Aut}(\mathbb{Q})$. Then f is generic if and only if $\mathcal{O}_f[\mathbb{Q}] \cong \mathbb{Q}$, and for each $\sigma \in \{+1, -1, 0\}$, the set of orbitals of parity σ is dense in $\mathcal{O}_f[\mathbb{Q}]$.

Question

Is an analogous statement true for **P**?

Answer

Partially.

Using orbitals to characterize generics of $\text{Aut}(\mathbf{P})$?

Theorem (I., 2020)

Using orbitals to characterize generics of $\text{Aut}(\mathbf{P})$?

Theorem (I., 2020)

If $f \in \text{Aut}(\mathbf{P})$ is generic,

Using orbitals to characterize generics of $\text{Aut}(\mathbf{P})$?

Theorem (I., 2020)

If $f \in \text{Aut}(\mathbf{P})$ is generic, then $(\mathcal{O}_f[\mathbf{P}], \langle \cdot \rangle_f^s) \cong \mathbf{P}$.

Using orbitals to characterize generics of $\text{Aut}(\mathbf{P})$?

Theorem (I., 2020)

If $f \in \text{Aut}(\mathbf{P})$ is generic, then $(\mathcal{O}_f[\mathbf{P}], \langle \cdot \rangle_f^s) \cong \mathbf{P}$. Moreover, the following sets are dense in $(\mathcal{O}_f[\mathbf{P}], \langle \cdot \rangle_f^s)$:

$$\{\mathcal{O}_f(x) : \text{par}(x, f) = +1\};$$

$$\{\mathcal{O}_f(x) : \text{par}(x, f) = -1\};$$

$$\{\mathcal{O}_f(x) : \text{par}(x, f) = 0, \text{sp}(x, f) = n\} \text{ for each } 1 \leq n \leq \infty.$$

Using orbitals to characterize generics of $\text{Aut}(\mathbf{P})$?

Theorem (I., 2020)

If $f \in \text{Aut}(\mathbf{P})$ is generic, then $(\mathcal{O}_f[\mathbf{P}], \langle \cdot \rangle_f^s) \cong \mathbf{P}$. Moreover, the following sets are dense in $(\mathcal{O}_f[\mathbf{P}], \langle \cdot \rangle_f^s)$:

$$\{\mathcal{O}_f(x) : \text{par}(x, f) = +1\};$$

$$\{\mathcal{O}_f(x) : \text{par}(x, f) = -1\};$$

$$\{\mathcal{O}_f(x) : \text{par}(x, f) = 0, \text{sp}(x, f) = n\} \text{ for each } 1 \leq n \leq \infty.$$

Remark

This is a *partial* answer to our goal because we do not know if the converse holds: whether this property implies genericity.

A different tactic

Note

Equip \mathbf{P} with a relation symbol for the graph of a unary function, and consider the resulting structures (\mathbf{P}, f) and (\mathbf{P}, g) for $f, g \in \text{Aut}(\mathbf{P})$.

A different tactic

Note

Equip \mathbf{P} with a relation symbol for the graph of a unary function, and consider the resulting structures (\mathbf{P}, f) and (\mathbf{P}, g) for $f, g \in \text{Aut}(\mathbf{P})$. Then f and g are conjugate iff $(\mathbf{P}, f) \cong (\mathbf{P}, g)$.

A different tactic

Note

Equip \mathbf{P} with a relation symbol for the graph of a unary function, and consider the resulting structures (\mathbf{P}, f) and (\mathbf{P}, g) for $f, g \in \text{Aut}(\mathbf{P})$. Then f and g are conjugate iff $(\mathbf{P}, f) \cong (\mathbf{P}, g)$.

Question

Can we study this structure to describe the conjugacy relation on $\text{Aut}(\mathbf{P})$?

A different tactic

Note

Equip \mathbf{P} with a relation symbol for the graph of a unary function, and consider the resulting structures (\mathbf{P}, f) and (\mathbf{P}, g) for $f, g \in \text{Aut}(\mathbf{P})$. Then f and g are conjugate iff $(\mathbf{P}, f) \cong (\mathbf{P}, g)$.

Question

Can we study this structure to describe the conjugacy relation on $\text{Aut}(\mathbf{P})$?

Answer

Yes and no.

A different tactic

Note

Equip \mathbf{P} with a relation symbol for the graph of a unary function, and consider the resulting structures (\mathbf{P}, f) and (\mathbf{P}, g) for $f, g \in \text{Aut}(\mathbf{P})$. Then f and g are conjugate iff $(\mathbf{P}, f) \cong (\mathbf{P}, g)$.

Question

Can we study this structure to describe the conjugacy relation on $\text{Aut}(\mathbf{P})$?

Answer

Yes and no. Finite substructures in this language don't "remember" enough.

A first-order language

Definition

A first-order language

Definition

- Let L be the language consisting of binary relations b_i , for $i \in \mathbb{Z}$.

A first-order language

Definition

- Let L be the language consisting of binary relations b_i , for $i \in \mathbb{Z}$.
- For each poset P and each $f \in \text{Aut}(P)$, let P_f be the L -structure obtained by letting:

$$b_i^f(x, y) \stackrel{\text{def}}{\iff} x \leq f^i(y).$$

A first-order language

Definition

- Let L be the language consisting of binary relations b_i , for $i \in \mathbb{Z}$.
- For each poset P and each $f \in \text{Aut}(P)$, let P_f be the L -structure obtained by letting:

$$b_i^f(x, y) \stackrel{\text{def}}{\iff} x \leq f^i(y).$$

- We identify $b_i^f(x, y)$ with its truth value in $\{0, 1\}$, and we consider the bi-infinite sequence $\mathbf{b}^f(x, y) \in 2^{\mathbb{Z}}$.

A first-order language

Definition

- Let L be the language consisting of binary relations b_i , for $i \in \mathbb{Z}$.
- For each poset P and each $f \in \text{Aut}(P)$, let P_f be the L -structure obtained by letting:

$$b_i^f(x, y) \stackrel{\text{def}}{\iff} x \leq f^i(y).$$

- We identify $b_i^f(x, y)$ with its truth value in $\{0, 1\}$, and we consider the bi-infinite sequence $\mathbf{b}^f(x, y) \in 2^{\mathbb{Z}}$.

Remark

$P_f \cong P_g$ iff f and g are conjugate in $\text{Aut}(P)$.

A first-order language

Definition

- Let L be the language consisting of binary relations b_i , for $i \in \mathbb{Z}$.
- For each poset P and each $f \in \text{Aut}(P)$, let P_f be the L -structure obtained by letting:

$$b_i^f(x, y) \stackrel{\text{def}}{\iff} x \leq f^i(y).$$

- We identify $b_i^f(x, y)$ with its truth value in $\{0, 1\}$, and we consider the bi-infinite sequence $\mathbf{b}^f(x, y) \in 2^{\mathbb{Z}}$.

Remark

$P_f \cong P_g$ iff f and g are conjugate in $\text{Aut}(P)$. But also, since L is infinite, finite substructures can encode a lot more information.

How to control \mathbf{b}^f sequences

Remark

Controlling behavior of \mathbf{b}^f sequences with finitary configurations is the name of the game here.

How to control \mathbf{b}^f sequences

Remark

Controlling behavior of \mathbf{b}^f sequences with finitary configurations is the name of the game here.

Example

Suppose the following is a substructure of \mathbf{P} :




How to control \mathbf{b}^f sequences

Remark

Controlling behavior of \mathbf{b}^f sequences with finitary configurations is the name of the game here.

Example

Suppose the following is a substructure of \mathbf{P} :



Let $p := \{(y, y'), (z, z')\} \in \text{Aut}_{<\omega}(\mathbf{P})$.

How to control \mathbf{b}^f sequences

Remark

Controlling behavior of \mathbf{b}^f sequences with finitary configurations is the name of the game here.

Example

Suppose the following is a substructure of \mathbf{P} :



Let $p := \{(y, y'), (z, z')\} \in \text{Aut}_{<\omega}(\mathbf{P})$. Then $\text{sp}(x, f) = \infty$ for every $f \in [p]$, and so $b_i^f(x, x) = 0$ for all $i \neq 0$.

What generic automorphisms of the random poset look like

Theorem (I., 2020)

What generic automorphisms of the random poset look like

Theorem (I., 2020)

Let $f \in \text{Aut}(\mathbf{P})$. Then f is generic iff the following hold:

What generic automorphisms of the random poset look like

Theorem (I., 2020)

Let $f \in \text{Aut}(\mathbf{P})$. Then f is generic iff the following hold:

(A) f has dense conjugacy class;

What generic automorphisms of the random poset look like

Theorem (I., 2020)

Let $f \in \text{Aut}(\mathbf{P})$. Then f is generic iff the following hold:

(A) f has dense conjugacy class;

(B) \mathbf{P}_f is ultrahomogeneous (as an L -structure);

What generic automorphisms of the random poset look like

Theorem (I., 2020)

Let $f \in \text{Aut}(\mathbf{P})$. Then f is generic iff the following hold:

- (A) f has dense conjugacy class;*
- (B) \mathbf{P}_f is ultrahomogeneous (as an L -structure);*
- (C) $\mathbf{b}^f(x, x)$ is eventually constant on both sides whenever $\text{par}(x, f) \neq 0$;*

What generic automorphisms of the random poset look like

Theorem (I., 2020)


Let $f \in \text{Aut}(\mathbf{P})$. Then f is generic iff the following hold:

- (A) f has dense conjugacy class;
- (B) \mathbf{P}_f is ultrahomogeneous (as an L -structure);
- (C) $\mathbf{b}^f(x, x)$ is eventually constant on both sides whenever $\text{par}(x, f) \neq 0$;
- (D) A technical condition — illustrated on the next slide — that forces $\mathbf{b}^f(x, y)$ to be eventually periodic on both sides for all $x, y \in \mathbf{P}$.

(...what's with condition (D)?)


Let $x, y \in \mathbf{P}$
such that
 $\text{sp}(x, f) = \infty$
and $y \notin f^{\mathbb{Z}}(x)$.

(...what's with condition (D)?)



Let $x, y \in \mathbf{P}$
such that
 $\text{sp}(x, f) = \infty$
and $y \notin f^{\mathbb{Z}}(x)$.
(D) asserts
there is some
chunk of $f^{\mathbb{Z}}(x)$
...

(...what's with condition (D)?)



Let $x, y \in \mathbf{P}$
such that
 $\text{sp}(x, f) = \infty$
and $y \notin f^{\mathbb{Z}}(x)$.
(D) asserts
there is some
chunk of $f^{\mathbb{Z}}(x)$
...
where this
configuration
exists.

Anything else?

Anything else?

Remark

We hoped model theory might be able to help, but there are some notable complications.

Anything else?

Remark

We hoped model theory might be able to help, but there are some notable complications.

Facts

Suppose $f \in \text{Aut}(\mathbf{P})$ is generic.

Anything else?

Remark

We hoped model theory might be able to help, but there are some notable complications.

Facts

Suppose $f \in \text{Aut}(\mathbf{P})$ is generic.

- The L -theory of \mathbf{P}_f is not ω -categorical.

Anything else?

Remark

We hoped model theory might be able to help, but there are some notable complications.

Facts

Suppose $f \in \text{Aut}(\mathbf{P})$ is generic.

- The L -theory of \mathbf{P}_f is not ω -categorical.
- \mathbf{P}_f is not ω -saturated.

Anything else?

Remark

We hoped model theory might be able to help, but there are some notable complications.

Facts

Suppose $f \in \text{Aut}(\mathbf{P})$ is generic.

- The L -theory of \mathbf{P}_f is not ω -categorical.
- \mathbf{P}_f is not ω -saturated.
- The relation $\text{sp}(x, f) = \infty$ is definable in \mathbf{P}_f , but not quantifier-freely.

Anything else?

Remark

We hoped model theory might be able to help, but there are some notable complications.

Facts

Suppose $f \in \text{Aut}(\mathbf{P})$ is generic.

- The L -theory of \mathbf{P}_f is not ω -categorical.
- \mathbf{P}_f is not ω -saturated.
- The relation $\text{sp}(x, f) = \infty$ is definable in \mathbf{P}_f , but not quantifier-freely. Thus, the L -theory of \mathbf{P}_f does not have QE.

Tack så mycket!