

## A NONLOCAL REACTION-DIFFUSION POPULATION MODEL WITH STAGE STRUCTURE

*Dedicated to Professor Paul Waltman on the occasion of his retirement*

DASHUN XU AND XIAO-QIANG ZHAO

**ABSTRACT.** A threshold dynamics and global attractivity of positive steady state are established in terms of principal eigenvalues for a nonlocal reaction-diffusion population model with stage structure, and the effects of spatial dispersal and maturation period on the evolutionary behavior are also discussed in two specific cases.

**1 Introduction** Recently, an increasing attention has been paid to nonlocal and time-delayed population models in order to study the effects of spatial diffusion and time delay on the evolutionary behavior of biological systems (see, e.g., [16, 13, 3, 1, 19, 17]). In the reality, species may drift from one spatial point at a time to another spatial point at another time, and may disperse from a domain to a larger domain. Moreover, the environment is often spatially heterogeneous. To describe the growth of a single species in a multi-patchy environment, certain delay differential equation models were proposed and analyzed in [12, 14, 20]. [13, 1, 3] also formulated the nonlocal and delayed reaction-diffusion models for a single species with stage structure, and established the existence of a family of traveling wave fronts for these models.

In order to obtain a general nonlocal and time delayed model for a single species in a bounded domain, we let  $u(t, a, x)$  be the density of individuals with age  $a$  at a point  $x$  at time  $t$ ,  $\tau$  be the length of the juvenile period. Denote by  $u_m(t, x)$  the density of mature adults. Then

---

Supported in part by the NSERC of Canada.

Keywords: Nonlocal and delayed model, persistence and extinction, positive steady states, global attractivity.

Copyright ©Applied Mathematics Institute, University of Alberta.